



Available Online at ESci Journals

Journal of Plant Breeding and Genetics

ISSN: 2305-297X (Online), 2308-121X (Print)
http://www.escijournals.net/JPBG



DISSECTING GENETIC DIVERSITY IN *GARCINIA XANTHOCHYMUS* USING ISSR AND RAPD MARKERS

^aJayesh S. Anerao, ^aVikas Jha, ^bLina Korgaonkar, ^cShanmuganan P. Devi, ^dNitin Desai

^aNational Facility for Biopharmaceuticals, G. N. Khalsa College, Matunga, Mumbai, Maharashtra, India.

^bDepartment of Bioinformatics, G. N. Khalsa College, Matunga, Mumbai, Maharashtra, India.

^cICAR Research Complex for Goa, Ela, Old Goa, India.

^dAmity University, Mumbai - Pune Expressway, Bhatan, Post – Somathne, Panvel, Mumbai, India.

ABSTRACT

Western Ghats of India is one of the megadiversity centres in the world. There are 15 species of *Garcinia* reported, of which many are endemic in nature. The genetic diversity in one of the same endemic species *Garcinia xanthochymus* were evaluated using ISSR and RAPD markers. A total of 15 individual plants from each location was analyzed. Six populations of *G. xanthochymus* were collected and screened using 16 ISSR and 10 RAPD markers. ISSR primers generated 80 bands of which 17 were polymorphic (21.25 % polymorphism) and RAPD primers produced 51 total bands of which 18 were polymorphic (35.29 % polymorphism). The ISSR and RAPD cluster analysis grouped them into two groups. The detected polymorphism is enough at species level and therefore the RAPD and ISSR can be used as efficient markers for genetic relatedness assessment in *G. xanthochymus*. The UPGMA and PCoA analysis distantly grouped these populations in two groups. The genetic diversity within the population is an important source for the evolution of the genera. *G. xanthochymus* has distinct populations in different geographical area.

Keywords: Genetic diversity, Genetic improvement, ISSR, Polymorphism, RAPD.

INTRODUCTION

The genus *Garcinia* L. belongs to the family Clusiaceae (Guttiferae). This family consists of approximately 200 species throughout the world, among which 36 species occur in India. The Western Ghats region is considered as a secondary centre of origin for *Garcinia* species, where six species are endemic to the Western Ghats only (Abraham *et al.*, 2006). As a genus, *Garcinia* contains many species which include *Garcinia indica* - *Garcinia indica* (Thouars) Choisy, *Garcinia gummigutta* - *Garcinia gummigutta* (L) Robson, *Garcinia hombroniana* - *Garcinia hombroniana* Pierre and *Garcinia xanthochymus* - *Garcinia xanthochymus* Hook. f. ex J. Anders that provide important resources in this region.

G. xanthochymus is distributed across the states of Maharashtra, Goa, Karnataka, Tamilnadu and Kerala. It is cultivated extensively in Southeast Asia where the fruit is used for preserves, jams, and curries. The dried fruit sap is called gamboge and provides a dye that is used in

watercolor paints. Fruits are antihelminthic and cardiotoxic; improves appetite. Ripe fruit is tonic, invigorating and alexipharmac; good in heart trouble and biliousness. Generally, sherbat made with "Amsul" (sun-dried slices of the fruit), with a little rock-salt, pepper, cumin, ginger and sugar, is administered in bilious conditions (Yusuf *et al.*, 1994). Xanthochymol present in the fruit is antibacterial against *Streptococcus faecalis* and *Klebsiella pneumoniae*; the action is better than tetracycline (Asolkar *et al.*, 1992). A new prenylated xanthone, 1, 3, 5, 6-tetrahydroxy-4, 7, 8-tri(3-methyl-2-butenyl) xanthone, was isolated from the wood of *G. xanthochymus* together with a known xanthone, garcinia xanthone. Their structures was determined by spectroscopic analysis. There are certain reports on the production of antibacterial silver nanoparticles using the endophytic bacterium *Bacillus cereus* isolated from *G. xanthochymus* (Sunkar and Nachiyar; 2012). However, the information on the genetic diversity of *G. xanthochymus* is not yet reported.

DNA fingerprinting is used to visualize DNA

* Corresponding Author:

Email: jayeshanerao@gmail.com

© 2016 ESci Journals Publishing. All rights reserved.

polymorphisms between samples. These fingerprints may be used as a tool for determining the identity of a specific DNA sample or to assess the relationship between samples. Fingerprints are also used as the source for genetic markers to generate linkage maps or to identify molecular markers linked to phenotypic traits and or genetic loci. Many DNA fingerprinting techniques have been developed in the past few years.

The genetic assessment of plant populations and species, for taxonomic, evolutionary, and ecological research, has been immensely assisted from the advancement of a range of molecular marker systems. Even though every molecular marker system is derived from distinctive principles, yet their function is to expose the genome-wide variability. Molecular markers offer a great advantage over the morphological and biochemical markers in species identification, phylogenetic studies and in mapping the genetic linkage (Suh *et al.*, 2011). Combined approach of both morphological and molecular markers was also carried out to identify and characterize the species. Different molecular markers have been used for the genetic diversity studies in different *Garcinia* species. The Phylogenetic analysis of some plants in the genus *Garcinia* using Internal Transcribed Spacer (ITS) sequence data and AFLP analysis have been studied by Chinawat Yapwattanaphun (2003). Molecular investigations using ITS were carried out for identifying the origin of Mangosteen (Nazre, 2014). Mansyah *et al.* (2013) evaluated the genetic diversity among and within *G. mangostana* L. (Mangosteen) trees. Polymorphism in *G. indica* was assessed by the morphological and molecular markers. Various populations collected from

Maharashtra screened with molecular markers, showed very less polymorphism (Thatte *et al.*, 2012).

The significance of the germplasm is recognized only when it is characterized (Rao *et al.*, 2002). Hence, the characterization of *G. xanthochymus* is necessary for its conservation and maintenance. The plants used in this study for characterization of *G. xanthochymus* were collected from different location of Western Ghats from Dapoli Maharashtra, ICAR Goa, Kadakeri, Siddhapur, Karnataka, GKVK Bangalore, Karnataka and NBPGR Thrissur, Kerala. Till date there are no reports based on the molecular markers for the genetic diversity for *G. xanthochymus*. Molecular markers are helpful in characterization of genetic divergence amongst several cultivars or populations of *G. xanthochymus* for the evaluation of genetic fidelity, identifying genes of commercial and agronomic interests, and enhancement via genetic transformation system. The aim of the present study was to fingerprint and explore the genetic relationship among the various populations of *G. xanthochymus* using ISSR and RAPD molecular markers.

MATERIAL & METHODS

Collection of Plant material: Fruits (Figure 1) and Seeds of *G. xanthochymus* were collected from various regions, Dapoli, (Maharashtra state), Indian Counsel of Agriculture and Research (Goa), Kadakeri, Siddhapur (Karnataka), Gandhi Krishi Vidyan Kendra, Bangalore (Karnataka) and National Bureau for Plant Germplasm Resources, Thrissur (NBPGR, Kerala) and planted in Khalsa college garden. The GPS locations and details of collection sites of samples are shown in Table 1 and Figure 2. Young leaves of *G. xanthochymus* were collected just prior to extraction of DNA from Khalsa college garden.



Figure 1. Fruits of *G. xanthochymus*



Figure 2. Map showing collection sites of *G. xanthochymus*

Table 1: List of the *G. xanthochymus* populations used in present study.

Accession No.	Location/Origin		GPS location	Sea Level
Population 1	Dapoli, Maharashtra		17° 45' 13.65" N	182m
Population 2	GKVK	Bangalore,	13° 4' 49.69" N	951m
	Karnataka		77° 34' 36.37" E	
Population 3	ICAR Complex Goa		15°30'52" N	58m
			73°55'01" E	
Population 4	Kadakeri,	Siddhapur,	14° 20' 27.3149" N	564m
	Karnataka		74° 53' 20.8465" E	
Population 5	NBPGR,	Thrissur,	10° 33' 5.32" N	33m
IC 552541-2	Kerala		76° 16' 30.60" E	
Population 6	NBPGR,	Thrissur	10° 33' 5.17" N	25m
IC 326809-1	Kerala		76° 16' 30.63" E	

The DNA was isolated from six populations of *G. xanthochymus*. About 3 grams of fresh leaflets were ground for DNA extraction. Total DNA was extracted and purified according to the slightly modified CTAB protocol (Doyle and Doyle 1987). Purified DNA concentrations were determined with electrophoresis on 0.8% agarose gel,

stained with 0.1 µg EtBr (ethidium bromide) and visualized using ultraviolet (UV) trans-illuminator. DNA was then used for PCR-amplification by using 16 ISSR and 10 RAPD primers obtained from the Eurofins. List of primers and their sequences were mentioned in Table 2 and Table 3.

Table 2: ISSR primers useful for the amplification of the six populations of *G. xanthochymus*.

Primer	Sequence from 5'-3'	Tm °C
Primer A1	ACACACACACACACTT	51°C
Primer A3	AGAGAGAGAGAGAGAGT	51°C
Primer A4	AGAGAGAGAGAGAGAGAA	51°C
Primer A5	AGAGAGAGAGAGAGAGTA	51°C
Primer A6	AGAGAGAGAGAGAGAGTT	51°C
17899B	CACACACACACAGG	49°C
ISSR14	GGAGTGTTGGTGGTG	53°C
ISSR11	AGAGAGAGAGAGAGAGGT	53°C
Primer 3c	CACACACACACACAAT	53°C
Primer 4c	CACACACACACACAAC	53°C
Primer 5c	CACACACACACACAGT	53°C
Primer 873c	GACAGACAGACAGACA	49°C
Primer M10	CACACACACACAAGG	49°C

PCR Reaction mixture: The chemicals were obtained from Thermo Fischer. Reactions were carried out in a total volume of 25 µl consisting of 100 ng of template DNA, 1X Taq. Buffer, 2.5 mM MgCl₂, 2.5 U Taq polymerase, 0.2mM dNTPs, and 0.5µM Primer. The total volume was made up using nuclease free water. Further, the reaction mixture was kept in the thermal cycler for amplification.

Amplification was performed under the following conditions: 5 min at 95°C for 1 cycle, followed by 1 min at 94°C, 1 min at annealing temperature (depending on the primer used), 1 min at 72°C for 45 cycles, and 7 min at

Table 3. RAPD primers useful for the amplification of the six populations of *G. xanthochymus*

Primer	Sequence from 5'-3'	Tm °C
RP 1	AAAGCTGCGG	32°C
RP 2	AACGCGTCGG	34°C
RP 3	AAGCGACCTG	32°C
RP 4	AATCGCGCTG	32°C
RP 5	AATCGGGCTG	32°C
RP 6	ACACACGCTG	32°C
RP 7	ACATCGCCCA	32°C
RP 8	ACCACCCACC	34°C
RP 9	ACCGCCTATG	32°C
RP 10	ACGATGAGCG	32°C

Note: Tm: Melting temperature.

72°C for final extension. PCR products were separated on 2% agarose gel stained with 0.1 µg/ml of ethidium bromide using 1X TAE buffer solution, and visualized under UV light.

Data Analysis: The amplified product of PCR was scored as presence of band (1) or absence of band (0) for the 90 population of *G.xanthochymus*. The relationship among the population was calculated through Jaccard coefficient. This analysis was performed using the un-weighted pair group method with arithmetic means (UPGMA) algorithm computed by DendroUPGMA (Garcia-Vallvé.S and

Puigbo.P,2002) and PAST version 3.12 (Paleontological Statistics).The PCoA analysis was carried out to visualize genetic diversity among population. The analysis was carried out using GenAlEx 6.5 (Peakall and Smouse, 2012).

RESULTS

ISSR analysis: Sixteen primers produced a total of 80 bands among the *G. xanthochymus* genotypes. The size of the amplified products ranged from 201 bp to 2000 bp. The number of scorable bands produced per primer ranged from 3 to 7. The total number of polymorphic bands and the percentage of polymorphism were 17 and 21.25 respectively (Table 4) (Figure 3 and 4).

ISSR A5 showed only one polymorphic band of 1300 bp, present in population of Dapoli, Bangalore and Goa (Figure 3) while in other three populations from Siddhapur, Thrissur 1(IC 552541-2) and Thrissur 2 (IC 326809-1) bands were absent. The ISSR trail of primer 3C showed three polymorphic bands in five populations whereas population of Thrissur1 showed absence of band in 900bp, 1500bp and 1600bp region (Figure 4).

Dendrogram was constructed from the data obtained from ISSR analysis. There were 2 clusters formed where first cluster consisted of population of Dapoli and Thrissur 2 while the rest of population lied in second cluster (Figure 7 A).

RAPD analysis: The total number of bands observed

among *G. xanthochymus* based on RAPD analysis was 51. The number of scorable bands produced per primer ranged from 3 to 7 and size of the products ranged from 300 bp to 1700 bp. The total number of polymorphic bands and the percentage of polymorphism were 18 and 35.29 respectively (Table 2) (Figure 5 and Figure 6).

Amplification produced by RP 4 primers, 1000 bp of band showing polymorphism in *G. xanthochymus*. It was produced in Dapoli and Siddhapur populations.

A comparison of the amplification pattern produced by RP 10 (Figure 6) showed six polymorphic bands of size 280 bp, 500 bp, 700bp, 800 bp and 900 bp in the six populations. 280 bp band was present in Dapoli, Thrissur 1 and Thrissur 2, 500 bp band was present in three populations Bangalore, Goa and Siddhapur respectively. 700 bp band was present in Bangalore, Goa, Siddhapur and Thrissur 2 populations while 800 and 900 bp bands were observed only in Bangalore, Goa and Siddhapur populations.

Dendrogram was constructed from the data obtained from RAPD analysis. There were 2 clusters formed where first cluster consisted of populations of Dapoli, Thrissur1 and Thrissur 2 while the rest of population lied in second cluster (Figure 7 B).

Table 4. Polymorphism observed in different populations of *G. xanthochymus* using molecular markers.

Markers	No. of primers	Total no. of bands	No. of polymorphic bands	% of polymorphism
RAPD	10	51	18	35.29%
ISSR	16	80	17	21.25%




Figure 3. ISSR fingerprints of six populations of *G. xanthochymus* generated with primer A5.




Figure 4. ISSR fingerprints of six populations of *G. xanthochymus* generated with primer 3C.




Figure 5. RAPD fingerprints of six populations of *G. xanthochymus* generated with primer RP4.




Figure 6: RAPD fingerprints of six populations of *G. xanthochymus* generated with primer RP 10.

Note: Loading Sequence: Lane 1-Ladder (L), 2-Dapoli (D), 3- Bangalore (B), 4- Goa (G), 5-Siddhapur (S), 6-Thrissur 1 (TR 1), 7- Thrissur 2 (TR 2).




Figure 7. UPGMA dendrogram based on the similarity matrix of *G. xanthochymus* obtained with the Jaccard coefficient for the data A) 16 ISSR primers B) 10 RAPD primers C) 26 RAPD+ISSR primers.

Cluster analysis: 126 bands were formed by analyzing the RAPD and ISSR techniques together which were used to construct a dendrogram based on Jaccard similarity coefficient. Cluster analysis performed using RAPD and ISSR generated a dendrogram that separated the genotypes into two distinct clusters (Figure 7 C). The first cluster included genotypes of Dapoli, Thrissur1 and Thrissur 2. It is divided into subcluster 1(a) consisting of Dapoli and Thrissur1 and Thrissur2 stands alone. The

second cluster included genotypes of Bangalore, Goa and Siddhapur. It is divided into subcluster 2(a) consisting Bangalore and Goa, and Siddhapur stands alone in cluster.

The Principal Coordinate Analysis performed with the complete set of molecular data (RAPD+ISSR) for 6 *G. xanthochymus* genotypes, accounted for 54.41%, 36.00%,9.58% of variation as explained by the first three

axes. Principal Coordinate 1 separates the genotypes belonging to cluster 2 from the cluster 1 (Figure 8).




Figure 8. Projection of the genotypes onto the plane defined by the Principal Coordinates 1 and 2, from the data of 26 markers (RAPD, ISSR).

DISCUSSION

The *Garcinia* species in India show rich source of biodiversity (Abraham *et al* 2006) and their conservation and utilization requires further characterization at inter and intra species level. Comparative studies on various *Garcinia* species involving RAPD, ISSR and SSR markers were successfully used by various researchers throughout the world (Mansyah *et al*, 2010; Qosim *et al.*, 2011; Mohan *et al*, 2012; Thatte *et al*, 2012; Tharachand *et al*, 2015).

The discriminative power of DNA markers used as tool to characterize the *Garcinia* species is essential because they can be used to assess the genetic diversity among *Garcinia* species. During the characterization of *G. xanthochymus* from Western Ghats, the discriminative power of RAPD and ISSR markers systems were evaluated. Among the two marker system employed 10 RAPD primers produced a total of 51 bands whereas 16 ISSR primers produced 80 bands. The level of polymorphism revealed by RAPD (35.29%) is higher than ISSR (21.25%). Thatte *et al.* (2012) also reported very less diversity in *G. indica* using morphological and molecular markers (RAPD and ISSR) collected from Maharashtra. The highest diversity within that population was 19.93%.

ISSR and RAPD markers can be used in population genetic studies of plant species as they effectively detect very low levels of genetic variation (Kumar *et al.*, 2014). They also may have potential for analyzing

biogeographic patterns among populations of a single plant species. In this study, we have shown that these markers revealed genetic variation among geographically separated samples of *G. xanthochymus* collected from different locations of Western Ghats. The results obtained in this study showed ISSR and RAPD markers measured sufficient polymorphism in *G. xanthochymus* and can be used in germplasm characterization and fingerprinting purposes.

Cluster analysis was carried out on two sets of marker profiling data based on combination of RAPD and ISSR. Cluster 1 is a heterogenous in terms of geographical location. The Thrissur1 and Thrissur2 genotypes were collected from Thrissur, Kerala which are placed along with Dapoli genotypes indicating that they share some genetic similarity.

Cluster 2 can also be called as heterogenous in terms of geographical location. Bangalore, Goa and Siddhapur populations were shown there can be a possible relationship among the three genotypes. This result was comparable to the genetic variation showed by UPGMA dendrogram. The same comparative results were found in UPGMA dendrogram and Principle Coordinate Analysis of *G. gummigutta* (Tharachand *et al* 2015). The Principle Coordinate Analysis (PCoA) plot generated by the RAPD primers showed the relationship of the populations and genetic variabilities in *G. gummigutta*.

This study may be indicative that due to geographical and topological difference the genotypes grouped into 2

clusters would have gone a significant variation that can be seen from their molecular characteristics.

Molecular markers ISSR and RAPD have been found useful for the study of genetic diversity of the present *G. xanthochymus* collection and numbers of markers analysed were sufficient to discriminate all genotypes. A better understanding of genetic diversity and its distribution is wanted to know for their conservation strategy. Of the markers studied RAPD showed high percentage of polymorphism than ISSR (35.29% and 21.25% respectively). Although the level of polymorphism revealed by ISSR and RAPD was sufficient to observed molecular insights and generate phylogenetic data in *G. xanthochymus*. Geographical conditions and surrounding environment made an impact on plant species which is evidenced by this study. Though the *G. xanthochymus* populations were not showing much diversity in their morphology, but showing enough genetic diversity at species level to discriminate the populations. These two molecular markers can be appropriate options since they are easier to implement and less costly. This study will help and encourage to other researchers and plant breeder to understand the molecular diversity of *G. xanthochymus*. The newest technologies in the field of DNA genotyping like DNA barcoding, Microarrays and Next Generation Sequencing (NGS) will efficiently work to the existing approaches of phylogenetic studies These approaches would be helpful not only in management of germplasm collections by avoiding redundancy but also in authentication of the genus *Garcinia*. Finally, the data obtained can be used for varietal survey and construction of germplasm collection and providing also additional information that could form the basis for the rational design of breeding programs.

ACKNOWLEDGEMENT

This work was supported by the Science and Engineering Research Board, Department of Science and Technology, New Delhi, Government of India, SERB Sanction Order No and date: SB/YS/LS-25/2014.

REFERENCES

- Abraham Z, Malik SK, Rao G, Narayanan and SL, Biju S. 2006. Collection and Characterisation of Malabar Tamarind [*Garcinia cambogia* (Gaertn.) Desr.]. *Genet Resour Crop Evol.*; 53(2): 401-406. <http://link.springer.com/article/10.1007/s10722-004-0584-y>
- Asolkar, L.V., K.K. Kakkar and O.J. Chakra, 1992. Second Supplement to Glossary of Indian Medicinal Plants with Active Principles Part-1 (A-K) 1965-1981. NISC, CSIR, New Delhi, India, pp: 265-266. <http://www.worldcat.org/title/glossary-of-indian-medicinal-plants-with-supplement/oclc/499428255>
- Doyle Doyle J.J, Doyle and J.L. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem Bull Bot Soc Amer* 19:11-15. <http://220.227.138.213/phytofura/phytoweb/protocols/IDoyle%20and%20Doyle%20octab%20method%20of%20dna%20isolation.pdf>
- Garcia-Vallvé.S and Puigbo.P, (2002). Horizontal gene transfer in glycosyl hydrolases inferred from codon usage in *Escherichia coli* and *Bacillus subtilis*. *Molecular Biology and Evolution* 16 (9): 1125-1134. <http://mbe.oxfordjournals.org/content/16/9/1125.full.pdf>
- Kumar, A., Mishra, P., Singh, S. C. (2014). Efficiency of ISSR and RAPD markers in genetic divergence analysis and conservation management of *Justicia adhatoda* L., a medicinal plant. *Plant Syst Evol* 300: 1409-1420. link.springer.com/article/10.1007/s00606-013-0970-z
- Mansyah E, Santoso PJ, Muas I, Sobir (2013). Evaluation of Genetic Diversity among and within Mangosteen (*Garcinia mangostana* L.) Trees. IV International Symposium on Tropical and Subtropical Fruits, *ISHS Acta Horticulturae*; 975: 73 -80. 10.17660/ActaHortic.2013.975.6
- Mohan S, Parthasarathy U, Babu KN (2012). Invitro and invivo adventitious bud differentiation from mature seeds of three *Garcinia* sps. *Ind J Nat Prod Resour* 3(1) 65-72 [http://nopr.niscair.res.in/bitstream/123456789/13812/1/IJNPR%203\(1\)%2065-72.pdf](http://nopr.niscair.res.in/bitstream/123456789/13812/1/IJNPR%203(1)%2065-72.pdf)
- Myers, N, Mittermeier RA, Mittermeier CG, Fonseca GAB and Kent J (2000). Biodiversity hotspots for conservation priorities. *Nature* 403: 853-858. <http://www.nature.com/nature/journal/v403/n6772/abs/403853a0.html>
- Nazre M (2014). New evidence on the origin of mangosteen (*Garcinia mangostana* L.) based on morphology and ITS sequence. *Genet Resour Crop Evol.*; 61: 1147-1158. 10.1007/s10722-014-0097-2

- Peakall and Smouse 2006 Peakall ROD, Smouse PE.. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol Ecol Notes* 6: 288-295. 10.1111/j.1471-8286.2005.01155
- Qosim WA, Patarapuwadol S, Watanabe KN (2011). Development of SSR markers of Mangosteen (*Garcinia mangostana* L.). *Int J Biotechnol* 2:1-8. <http://www.interestjournals.org/IRJOB>
- Rao VR, Hodgkin T (2011). Genetic diversity and conservation and utilization of plant genetic resources. *Plant Cell, Tissue and Organ Culture* 68(1): 1-19. <https://www.cbd.int/doc/articles/2002-/A-00109.pdf>
- Suh JK, Hettterscheid W, Lee AK, Hong J, Roh MS (2011) Identification and evaluation of *Forsythia* germplasm using molecular markers. *Genet Resour Crop Evol.* 58: 1225-1235. 10.1007/s10722-010-9656-3
- Sunkar S. and Nachiyar CV (2012). Biogenesis of antibacterial silver nanoparticles using the endophytic bacterium *Bacillus cereus* isolated from *Garcinia xanthochymus* *Asian Pac. J Trop Biomed* 2(12): 953-959. 10.1016/S2221-1691(13)60006-4
- Tharachand C1, Immanuel Selvaraj C and Abraham Z (2015). Molecular Insights into the Genetic Diversity of *Garcinia cambogia* Germplasm Accessions. *Braz Arch Biol Technol* 58(5)765-772. <http://dx.doi.org/10.1590/S1516-89132015050197>
- Thatte KS, Khandekar RG, Deodhar MA (2012). Assessment of diversity in *Garcinia indica* (Dupetit-Thouars.) Choisy. using morphological and molecular markers. *Journal of tropical Agriculture* 50: 30-36. <http://www.jtropag.in/index.php/ojs/article/viewFile/1091/300>
- Yapwattanaphun C. (2003) Dissertation: Phylogenetic analysis of some plants in the Genus *Garcinia* using ITS sequence data and AFLP analysis. A Dissertation Submitted in Partial Fulfillment of the Requirements for the Degree of Doctor of Philosophy (Horticulture) Graduate School, Kasetsart University 2003 ISBN 974-359-434-5.
- Yusuf, M., Wahab M.A. , Chowdhury J.U. and Jaripa Begum (1994). Medicinal plants of Bangladesh. BCSIR Laboratories, Bangladesh. <http://www.mpbd.info/related-link.php>.